arXiv:1304.6425v2 [quant-ph] 27 May 2013

Lack of measurement independence can simulate quantum correlation even when signaling cannot

Manik Banik[1, *]

[1] *Physics and Applied Mathematics Unit, Indian Statistical Institute, 203 B.T. Road, Kolkata-700108, India*

In a world described by quantum mechanics, one cannot have determinism, no-signaling and free will all together. Under these three assumptions the set of statistical correlations are restricted to the set of Bell local correlations. In standard Bell scenario any non-local correlation has a deterministic explanation if one either sacrifices free will or allows signaling. Recently in [Phys. Rev. Lett. **108**, 200401 (2012)] F. Buscemi generalizes the standard Bell scenario into 'semi-quantum' Bell scenario. Interestingly, in 'semi-quantum' Bell game scenario we show existence of correlations that cannot be obtained by sacrificing no-signaling by any amount, whereas such correlations can be obtained by giving up a small degree of free will.

PACS numbers: 03.67.Mn, 03.65.Ud

In the early days of quantum mechanics, Einstein and his colleagues designed a thought experiment [1] intended to reveal what they believed to be inadequacies of quantum mechanics and they have dreamt a causal world where no intrinsic randomness is present. More than three decades later Bell showed that if an experimenter enjoys total free will then quantum mechanics is indeed in contradiction with local realism. Under validation of three different assumptions, namely, determinism, no-signaling and free will, an inequality can be derived, famously known as Bell-CHSH inequality [2, 3], which is violated by quantum correlations and the experimental verification of quantum violation is showed in [4]. Correlations that violates Bell inequality are called non-local correlations and the amount of nonlocality is quantified by the amount of Bell inequality violation [2, 3]. Interestingly, sacrificing no-signaling or experimental free will one can obtain any amount of bell inequality violation [5]. Particularly, for non-local quantum correlations there exists various interesting simulation protocols sacrificing determinism [6], sacrificing no-signaling [7, 8] as well as sacrificing free will [9, 10]. In [5, 11, 12], various interesting relations have been derived showing interrelations among the degree of weakening of various assumptions to simulate a given Bell non-local correlation.

As any Bell non-local correlation can be explained by giving up no-signaling assumption as well as giving up free will assumption in suitable amount, therefore in standard Bell scenario one can not make an *operational* distinction between strength of these two different assumptions. In this letter we show that modification of standard Bell game scenario to 'semi-quantum' Bell game scenario [14] allows one to make such *operational* distinction. More explicitly said, we show that in 'semi-quantum' Bell game one can get correlations that cannot be obtained by local operation and classical communication even if there is no restriction on the amount of classical communication; whereas such correlations can be obtained by sacrificing a small degree of free will. Thus 'semi-quantum' game provides a novel framework where one can make *operational* distinction between two very different assumptions, no-signaling and free will. We also argued that like in standard Bell scenario, such operational distinction between no-signalling and measurement independence is not possible in 'quantum refereed steering games? scenario, introduced by Cavalcanti *et.al.* in [15].

The standard Bell game scenario [2, 3] involves two spatially separated players, say Alice and Bob. A referee picks indices $s \in \mathcal{S}$ and $t \in \mathcal{T}$ at random and sends them separately to Alice and Bob, respectively. The two players must compute answers $x \in \mathcal{X}$ and $y \in \mathcal{Y}$, respectively, and send them to the referee, who will then pay them both. First, the players are told the rules of the game. Knowing the rules, the players are allowed to agree on any strategy and to share any variable $\lambda \in \Lambda$. The distribution $\rho(\lambda)$ of the shared variable λ must be independent of the referee's questions, say s and t; otherwise the free will assumption will not be satisfied. After starting the game the two players are forbidden to communicate with each other. when each of s, t, x, y are binary, the canonical no-go result is represented by the Bell-CHSH inequality, which puts constraint that have to be satisfied by Bell-local correlations. Alice and Bob can obtain a better than classical payoff by sharing bipartite entangled quantum states. Though all bipartite pure entangle states give better than classical payoff [16], there exist mixed entangled states for which one can never obtain better than classical payoff [17]. In this sense nonlocality [18] and entanglement [19] are two distinct concepts in quantum mechanics.

Very recently, F. Buscemi [14] generalizes the standard Bell non-local game into 'semi-quantum' non-local game. In 'semi-quantum' non-local game referee picks indices $s \in \mathcal{S}$ and $t \in \mathcal{T}$ at random and sends quantum states τ^s and ω^t to Alice and Bob, respectively, without reveling the classical indices s and t. In other word, in standard non-local game the referee is allowed to ask only *classical questions*, but in 'semi-quantum' non-local game the referee is given the power to ask *quantum questions.* In this case, also, the free will assumption is applied i.e., the distribution $\rho(\lambda)$ of the shared variable λ is independent of referee's *quantum questions* and the constraint of no communication between the two players must satisfies. An immediate consequence of Buscemi's 'semi-quantum' non-local game framework is that given any entangled state there exists a 'semi-quantum' non-local game for which the entangled state gives better payoff than achievable by sharing any separable state.

---

*Electronic address: manik11ju@gmail.com

In [15], Cavalcanti *et. al.* discuss the work of Buscemi, and offer a perspective on the operational significance of that work. They have also made a non-trivial extension of Buscemi's work by introducing 'quantum refereed steering games' and showing that all steerable states exhibit correlations in 'quantum refereed steering games' that can not be achieved by *local operation with steering and shared randomness* (LOSSR). Using Buscemi's 'semi-quantum' game framework, Branciard *et.al.* introduce the concept of Measurement-Device-Independent Entanglement Witnesses, which allow one to demonstrate entanglement of all entangled quantum states with untrusted measurement apparatuses [20]. In [21], Rosset *et.al.* discussed about *quantum input simulation task*, and proved the no-go result regarding simulation of these correlations with local operation and classical communication (LOCC) [22]. This surprising result is possible because in *quantum input simulation task* referee sends nonorthogonal quantum states to Alice and Bob, who can not discriminate the states even if they are allowed to communicate with each other.

In this letter, we show that sacrificing free will one can simulation the correlations achieved in *canonical quantum input simulation task.* To present our work we first describe *canonical quantum input simulation task*, introduced in [21] and explained the no-go result regarding simulation of these correlations with LOCC. We then formulate an explicit simulation protocol for such correlations by sacrificing the free will. Thus in 'semi-quantum' game framework one can make an operational distinction between lack of free-will and signaling. We also pointed out that in *quantum refereed steering games*, introduced in [15], one cannot obtain such operational distinction as we have constructed an explicit simulation protocol for such games with finite communications.

The scenario of *quantum-input simulation task* (QIST) [21] is defined via the 5-tuple $(\varrho_{AB}, \{\tau^s_{A'}\}, \{\omega^t_{B'}\}, \{\mathcal{A}^x_{A'A}\}, \{\mathcal{B}^y_{BB'}\})$; where ϱ_{AB} is the shared entangled state between two player Alice and Bob, $\{\tau^s_{A'}\}$ and $\{\omega^t_{B'}\}$ are quantum inputs of referee to Alice and Bob respectively, $\{\mathcal{A}^x_{A'A}\}$ and $\{\mathcal{B}^y_{BB'}\}$ are POVM [23] elements acting respectively on the systems $A'A$ and BB'. To define *canonical quantum-input simulation task*, introduce in [21], let us consider a state ϱ_{AB} of dimension $d_A \times d_B$, and Alice and Bob perform local measurements $\mathcal{A}^x_{A'A}$ and $\mathcal{B}^y_{BB'}$ with binary outcomes $x, y \in \{1, 0\}$; where the outcome x or $y = 1$ corresponds to the successful projection onto the maximally entangled state $|\phi^+_d\rangle = \Sigma^{d-1}_{k=0}|kk\rangle/\sqrt{d}$. Thus

$$\mathcal{A}^1_{A'A} = |\phi^+_{d_A}\rangle\langle\phi^+_{d_A}| \ , \qquad \mathcal{A}^0_{A'A} = \mathbb{I} - |\phi^+_{d_A}\rangle\langle\phi^+_{d_A}| \tag{1}$$

$$\mathcal{B}^1_{BB'} = |\phi^+_{d_B}\rangle\langle\phi^+_{d_B}| \ , \qquad \mathcal{B}^0_{BB'} = \mathbb{I} - |\phi^+_{d_B}\rangle\langle\phi^+_{d_B}| \tag{2}$$

The input states are chosen to have the same dimensions as the respective subsystem of A and B in ϱ_{AB}, $|\tau^s\rangle \in \mathbb{C}^{d_A}$, $|\omega^t\rangle \in \mathbb{C}^{d_B}$, and constructed such that the corresponding density matrices $\{|\tau^s\rangle\langle\tau^s|\}$, $\{|\omega^t\rangle\langle\omega^t|\}$ span the space of linear operators acting on $\mathbb{C}^{d_A}$ and $\mathbb{C}^{d_B}$ respectively. As a concrete example of *canonical quantum-input simulation task* Rosset *et. al.* allow Alice and Bob to share singlet states $\varrho_{AB} = |\psi^-\rangle\langle\psi^-|$. The referee chooses his *quantum questions* from the vertices of a regular tetrahedron on the surface of the Bloch sphere. Using $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ the vector of Pauli matrices, and $\vec{v_1} = \frac{(1,1,1)}{\sqrt(3)}$, $\vec{v_2} = \frac{(1,-1,-1)}{\sqrt(3)}$, $\vec{v_3} = \frac{(-1,1,-1)}{\sqrt(3)}$, $\vec{v_4} = \frac{(-1,-1,1)}{\sqrt(3)}$, we denote, for $x, y = 1, 2, 3, 4$:

$$|\tau^x\rangle\langle\tau^x| = \frac{\mathbb{I} + \vec{v_x}.\vec{\sigma}}{2} \ , \quad |\omega^y\rangle\langle\omega^y| = \frac{\mathbb{I} + \vec{v_y}.\vec{\sigma}}{2} \tag{3}$$

The resulting correlations are then:

$$\begin{aligned}
P_{|\psi^-\rangle}(0,0||\tau^s\rangle, |\omega^t\rangle) &= \begin{cases} \frac{1}{2} & \text{if s=t} \\ \frac{7}{12} & \text{otherwise} \end{cases} \\
P_{|\psi^-\rangle}(0,1||\tau^s\rangle, |\omega^t\rangle) &= \begin{cases} \frac{1}{4} & \text{if s=t} \\ \frac{1}{6} & \text{otherwise} \end{cases} \\
P_{|\psi^-\rangle}(1,0||\tau^s\rangle, |\omega^t\rangle) &= \begin{cases} \frac{1}{4} & \text{if s=t} \\ \frac{1}{6} & \text{otherwise} \end{cases} \\
P_{|\psi^-\rangle}(1,1||\tau^s\rangle, |\omega^t\rangle) &= \begin{cases} 0 & \text{if s=t} \\ \frac{1}{12} & \text{otherwise} \end{cases}
\end{aligned} \tag{4}$$

Rosset *et. al.* then pointed out that, if instead of singlet state Alice and Bob are given pre-shared variable λ (of course the distribution of the variable λ is independent of referee's *quantum question*) and are allowed to communicate to each other, still they cannot reproduce the desired correlation (4). Rosset *et. al.*then extended this no-go result in the sense that any entangled state can generate correlations in a QIST that cannot be simulated by local operation assisted by classical communication, even if there is no limitation on classical communication (Corollary of the Theorem 1 in [21]). Thus in this generalized 'semi-quantum' scenario relaxation of no-signaling assumption has no use to simulate the correlation achieved from entangled states.. They actually got such a powerful no-go result as LOCC between Alice and Bob do no help in unambiguous discrimination of the nonorthogonal (and moreover linearly dependent) states sent to them by the referee. Interestingly, we will show that if Alice and Bob are allowed to share classical variable whose distribution depends on the *quantum input* of the referee then the correlation (4) can be reproduced even if Alice and Bob are not allowed to communicate and of course they don't need to discriminate the quantum states sent to them.

Let Alice and Bob share hidden variable Λ, taking 4 distinct values $\lambda = \lambda_1, \lambda_2, \lambda_3, \lambda_4 \in \Lambda$. These variables are distributed according to one of two probability distributions $p_{\_Sa(\lambda)}$ and $p_{\_Di(\lambda)}$:
for $s = t$

$$\{p_{\_Sa(\lambda)}\} \equiv \{p(\lambda|s,t)\} := \{1/2, 1/4, 1/4, 0\} \qquad (5)$$

and for $s \neq t$

$$\{p_{\_Di(\lambda)}\} \equiv \{p(\lambda|s,t)\} := \{7/12, 1/6, 1/6, 1/12\} \qquad (6)$$

The outcomes of Alice and Bob are fully determined by the parameter Λ as

$$x = f(\lambda), \qquad y = g(\lambda) \qquad (7)$$

where,

$$\begin{aligned} f(\lambda_1) = f(\lambda_2) = 0, \quad f(\lambda_3) = f(\lambda_4) = 1 \\ g(\lambda_1) = g(\lambda_3) = 0, \quad g(\lambda_2) = g(\lambda_4) = 1 \end{aligned}$$

The corresponding correlations are of the form

$$\begin{aligned} p(x,y|s,t) &= \sum_{\lambda} p(x,y|x,t,\lambda) p(\lambda|s,t) \\ &= \sum_{\lambda} \delta_{x,f(\lambda)} \delta_{y,g(\lambda)} p(\lambda|s,t) \qquad (8) \end{aligned}$$

where $\delta_{0,0} = \delta_{1,1} = 1$ and $\delta_{0,1} = \delta_{1,0} = 0$. It is clear that correlation (8) is same as correlation (4). It is straight forward to construct a model with lack of free will for any two qubits entangled state by using the probability distribution generated from the entangled state and quantum questions suggested above.

Now coming to the question of how much free will one has to sacrifice to reproduce correlation (4), we follow the quantification scheme of Hall, introduced in [5, 9]. Measurement independence is the property that the distribution of the underlying variable is independent of the measurement settings or referee's input question, i.e.,

$$p(\lambda|s,t) = p(\lambda|s',t') \qquad (9)$$

The degree to which measurement independence is violated can most simply quantified by the variational distance

$$M := \sup_{s,s',t,t'} \sum_{\lambda \in \Lambda} |p(\lambda|s,t) - p(\lambda|s',t')| \qquad (10)$$

Clearly $0 \leq M \leq 2$, where two extreme values signify complete measurement independence and complete measurement dependence, respectively. The fraction of measurement independence or free will is defined by

$$F := 1 - M/2 \qquad (11)$$

Thus, $0 \leq F \leq 1$, with $F = 0$ corresponding the case where no experimental free will can be enjoined and $F = 1$ corresponding the case with complete experimental free will. In [9] a local deterministic model for projective spin correlations for singlet state has been presented via giving up experimental free will (or measurement independence) by 14%.

For the above reduced free will model of *canonical quantum input simulation task* for singlet we have

$$M = |\frac{7}{12} - \frac{1}{2}| + 2|\frac{1}{6} - \frac{1}{4}| + |\frac{1}{12} - 0| = \frac{1}{3}$$

and the amount of experimental free will is given by

$$F = 1 - \frac{1}{2.3} \cong 83.3\%$$

The degree of measurement dependence of the model can also be quantified by the mutual information between Λ and the labels sent by the referee, i.e., $H(\Lambda : \mathcal{S}, \mathcal{T})$ [5, 13]. This depends on the probability distribution $p(s,t)$, however for $p(s,t) = 1/16$ one has $p(s,t,\lambda) = p(\lambda|s,t)p(s,t) = \frac{1}{16}p(\lambda|s,t)$, which yields $p(\lambda) = \frac{1}{4}p_{\_Sa(\lambda)} + \frac{3}{4}p_{\_Di(\lambda)}$. Mutual information is calculated by the following formula:

$$H(\Lambda : \mathcal{S}, \mathcal{T}) = \sum_{s,t,\lambda} p(s,t,\lambda) \log_2 \frac{p(s,t,\lambda)}{p(s,t)p(\lambda)} \qquad (12)$$

For the above model it becomes $H(\Lambda : \mathcal{S}, \mathcal{T}) \simeq 0.03705$. The *measurement dependence capacity* [9] of a given model may be defined by:

$$\mathcal{C}_{\text{meas dep}} := \sup_{p(s,t)} H(\mathcal{S}, \mathcal{T} : \Lambda) \qquad (13)$$

which turns out to be 0.05778 bits for our model (see [25]).

It is important to notice that distribution of the variable Λ depends on referee's input in such a way (Eqn.(5) and Eqn.(6)) that having accesses the variable Λ neither Alice nor Bob can guess the classical indices of their quantum states. As for example, if there shared variable is λ_4, then they can conclude that there indices are not same i.e. $s \neq t$; but noway they can guess their individual indices s and t respectively. Our model also makes clear that Rosset *et al.* 'no-go' result have been derived under "free will" assumption, which assumes no correlation between the referee's labels (s,t) and any other physical parameter Λ that propagates to the future (and hence to Alice and Bob). This is a very strong assumption, and our model shows that even a small lack of free will allows simulation of the correlations by non-quantum means without any need of communication. It is also interesting to study whether such simulation protocol with reduced free will exist for all 'semi-quantum' non-local games. Our intuition go affirmative in this case. Answering the above question positively one may also be interested in constructing a general simulation protocol with lack of free will for such ‘semi-quantum’ non-local games.

Besides non-locality quantum correlation exhibits another distinct non classical feature called steering, first introduced by Schrodinger [26] and recently by Wiseman and his collaborators [27, 28]. In [15], Cavalcanti *et. al.* have made an nontrivial extension of Buscemi’s work. They introduces *quantum refereed steering game* and shows that all steerable states exhibit correlations in *quantum refereed steering game* that can not be simulated by *local operation with steering and shared randomness* (LOSSR) maps [15] as well as by LOSR maps [24]. In the context of this paper it is interesting to ask the question whether *quantum refereed steering game* can make such operational distinction between free will and no-signaling. We answer this question in negative, as it can been shown that any correlation achieved in *quantum refereed steering game* can be simulated with the help of LOCC resources (see [25]).

Experimental free will and no-signaling are two very popular physically motivated assumptions. Various important no-go results in quantum foundation have been derived using these assumptions. While Bell’s theorem [2] and recently proved Colebeck and Renner’s result [29] uses both the assumption, Kochen-Specker no-go result [30] uses only free will assumption among these two. In an interesting discussion [31] of the measurement problem for a PR-box, J. Bub pointed out that, if Bob has access to a parameter or ontic state λ that completely specifies the output values for both of his possible inputs, or even specifies whether these output values are the same or different, then either there is a violation of no-signaling principle or Alice’s choice of input is not free but depends on the value of λ. Actually Bub’s conclusion holds for any standard Bell non-local correlation. Interestingly our result shows that in ‘semi-quantum’ non-local game there exists correlation whose deterministic explanation is possible only if free-will assumption is relaxed. Therefore in ‘semi-quantum’ non-local game scenario our result suggests further study of simulation protocol with reduced free-will.

I. MEASUREMENT DEPENDENCE CAPACITY

The *measurement dependence capacity* [9] of a given model may be defined by maximizing the mutual information over all possible distributions of measurement settings:

$$\mathcal{C}_{\text{meas dep}} := \sup_{p(s,t)} H(\mathcal{S},\mathcal{T}:\mathcal{V})$$

It is straightforward to find the maximum mutual information required to implement our model, over all possible $p(s,t)$, given that the hidden variable distribution depends only on whether $s=t$ or otherwise. Let

$$P := \sum_{s=t} p(s,t)$$

be the probability of sending the same label to Alice and Bob. Then, the mutual information is given by the simple formula

$$\begin{aligned} H(\mathcal{S},\mathcal{T}:\mathcal{V}) = {} & H[P p_{\mathcal{S}(v)} + (1-P) p_{\mathcal{D}(v)}] \\ & -PH[p_{\mathcal{S}(v)}] - (1-P)H[p_{\mathcal{D}(v)}], \end{aligned}$$

where $H[p]$ denotes the Shannon entropy of distribution p. It corresponds to a communication channel where one of two signals $s=t$ and $s\neq t$ are sent with probabilities P and $1-P$, and $\mathcal{V}$ is used to distinguish between them. Thus,the mutual information only depends on the probability P, and the maximum can easily be found numerically (see Fig(1))by using MATHEMATICA.



FIG. 1: This is $H(\mathcal{S},\mathcal{T}:\mathcal{V})$ Vs P plot. The point $A\cong(0.593, 0.057)$ correspond to maximum mutual information i.e., *measurement dependence capacity* of the model; and the point $B\cong(0.25, 0.037)$ correspond to mutual information between the variable $\mathcal{V}$ and $(\mathcal{S},\mathcal{T})$ when $p(s,t)$ is uniform.

II. SIMULATION OF *QUANTUM REFEREED STEERING GAME* BY LOCC RESOURCES

In *quantum refereed steering game*, a referee picks classical induces $s\in\mathcal{S}$ and $t\in\mathcal{T}$ at random, and sends the orthogonal quantum states $\pi^s\in\mathcal{H}_{A_0}$ to one party (say Alice) and sends quantum states $\omega^t\in\mathcal{H}_{B_0}$ (in general non orthogonal) to Bob, without revealing the actual induces s and t though. As the states π^s’s are orthogonal, it is required that $\dim(\mathcal{H}_{A_0})\geq\mathfrak{C}(\mathcal{S})$, where $\mathfrak{C}(\mathcal{S})$ denote the cardinality of the set $\mathcal{S}$. Since Alice receives orthogonal states, it is possible for her to perfectly determine the classical index s. On the other hand if ω^t’s are chosen from linearly dependent set then it is impossible for Bob to determine the classical index t. Now given the bipartite state ρ_{AB} shared between Alice and Bob, let us denote the set of all joint quantum conditional probabilities by $\mu(\rho_{AB}) := \{\mu(x,y|s,t) \mid x\in\mathcal{X};\ y\in\mathcal{Y};\ s\in\mathcal{S};\ t\in\mathcal{T}\}$, where $x\in\mathcal{X}$ and $y\in\mathcal{Y}$ are classical results of Alice and Bob respectively that has to convey to the referee. $\mu(x,y|s,t)$ is computed as:

$$\mu(x,y|s,t) = Tr[(P^x_{A_0A}\otimes Q^y_{BB_0})(\pi^s_{A_0}\otimes\rho_{AB}\otimes\omega^t_{B_0})],$$

where $P \equiv (P^x_{A_0A}; x \in \mathcal{X} \mid \sum_{x\in\mathcal{X}} P^x_{A_0A} = \mathbf{1}) \in \mathcal{M}(A_0A;\mathcal{X})$ and $Q \equiv (Q^y_{B_0B}; y \in \mathcal{Y} \mid \sum_{y\in\mathcal{Y}} Q^y_{B_0B} = \mathbf{1}) \in \mathcal{M}(BB_0;\mathcal{Y})$. By $\mathcal{M}(A;\mathcal{X})$, we denote the convex set of all $\mathcal{X}$-POVMs on A.

By simulation of the correlations $\mu(\rho_{AB})$, we mean that Alice and Bob are not allowed to share the state ρ_{AB}, but they have to reproduce the correlations in a given operational paradigm like LOSR or LOSSR or LOCC. Now if the state ρ_{AB} is steerable then according to the result of Cavalcanti *et.al* [15] we know that $\mu(\rho_{AB})$ cannot be simulated by LOSSR resources and therefore by LOSR resources. Interestingly if there allowed operational paradigm is LOCC then they will be succeed in the simulation task.

LOCC simulation protocol

Step 1: Bob locally prepares the state ρ_{AB} in his laboratory and takes another particle described by the Hilbert space $\mathcal{H}'_B \cong \mathcal{H}_{A_0}$.

Step 2: Receiving orthogonal quantum states π^s's from referee, Alice reveals the classical indices s by performing Von-Newman measurement in $\{\pi^s\}_{s\in\mathcal{S}}$ basis.

Step 3: Alice communicates her indices to Bob by using $\log_2 \mathfrak{C}(\mathcal{S})$ bits, where $\mathfrak{C}(\mathcal{S})$ be the cardinality of the set $\mathcal{S}$.

Step 4: Receiving the information of Alice's index s, Bob prepares the system B' in the state π^s.

Step 5: Receiving the quantum states ω^t's from referee, Bob performs the required POVM $P \in \mathcal{M}(A_0A;\mathcal{X})$ and $Q \in \mathcal{M}(BB_0;\mathcal{Y})$ and reproduces the joint conditional probability distributions $\mu(x, y|s, t)$.

Step 6: Bob communicates the result x to Alice by using $\log_2 \mathfrak{C}(\mathcal{X})$ bits.

Note that, if the condition of the simulation task demands that Alice and Bob have to just produce the correlations then the process ends at **Step 5** and they will be succeed in the simulation task by means of one way communications. But it the condition of the task further demands that the outcomes $x \in \mathcal{X}$ and $y \in \mathcal{Y}$ haves to be supply to referee by the respective parties, then they have to follow all the six steps and in this case they need both way communications.

Of course, Bob has to have the resources to be able to prepare a suitable bipartite state ρ_{AB} that exhibits steering. But this might be particularly relevant in a scenario where Alice and Bob want to pretend to be able to share such a state over long distances. In this scenario, they can reproduce the correlation by not sharing the state at all, but having Bob keeping it with him. Actually by the above clever means, they will be able to preserve sufficient entanglement shared between them. Thus following the above protocol all *quantum referred steering correlations* can be simulated in LOCC operational framework with bounded amount of classical communications.

Acknowledgement: The author acknowledges useful discussions with A. Rai, S. Das, Md R. Gazi, S. Kunkri and is grateful to G. Kar for discussion about elegant LOCC simulation protocol for quantum refereed steering correlations, and also gratefully acknowledges Michael J.W. Hall for suggesting a simple version of the reduced free will model. It is also my pleasure to thank F. Buscemi for useful discussion about his work [14].

[1] A. Einstein, B. Podolsky, and N. Rosen,Phys. Rev. **47**, 777 (1935).
[2] J. S. Bell, Physics **1**, 195 (1964).
[3] J. F. Clauser, M.A. Horne, A. Shimony and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
[4] A. Aspect, P. Grangier, and G. Roger, Phys. Rev. Lett. **47**, 460 (1981); A. Aspect, P. Grangier, and G. Roger, Phys. Rev. Lett. **49**, 91 (1982); A.Aspect, J. Dalibard, and G. Roger, Phys. Rev. Lett. **49**, 1804 (1982).
[5] M.J.W. Hall, Phys. Rev. A **84**, 022102 (2011).
[6] N.J. Cerf, N. Gisin, S. Massar, and S. Popescu, Phys. Rev. Lett. **94**, 220403 (2005).
[7] B.F. Toner and D. Bacon, Phys. Rev. Lett. **91**, 187904 (2003).
[8] S. Massar, D. Bacon, N. Cerf, and R. Cleve, Phys. Rev.A **63**, 052305 (2001); J. Degorre, S. Laplante, and J. Roland, Phys. Rev. A **72**, 062314 (2005).
[9] M.J.W. Hall, Phys. Rev. Lett. **105**, 250404 (2010).
[10] M. Banik, MD R. Gazi, S. Das, A. Rai, S. Kunkri, J. Phys. A: Math. Theor. **45**, 205301 (2012).
[11] M.J.W. Hall, Phys. Rev. A **82**, 062117 (2010).
[12] G. Kar, MD. R. Gazi, M. Banik, S. Das, A. Rai, S. Kunkri, J. Phys. A: Math. Theor. **44**, 152002 (2011).
[13] J. Barrett, N. Gisin, Phys. Rev. Lett. **106**, 100406, (2011).
[14] F. Buscemi, Phys. Rev. Lett. **108**, 200401 (2012).
[15] E. G. Cavalcanti, M. J. W. Hall, and H. M. Wiseman, Phys. Rev. A **87**, 032306 (2013).
[16] N. Gisin, Phys. Lett. A **154**, 201 (1991).
[17] R. F. Werner, Phys. Rev. A **40**, 4277 (1989); J. Barrett, Phys. Rev. A **65**, 042302 (2002).
[18] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, S. Wehner, arXiv[quant-ph] : 1303.2849; and reference therein.
[19] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Rev. Mod. Phys. **81**, 865 (2009); and reference therein.
[20] C. Branciard, D. Rosset, Y.C. Liang, and N. Gisin, Phys. Rev. Lett. **110**, 060405 (2013).
[21] D. Rosset, C. Branciard, N. Gisin, and Y. C. Liang, arXiv[quant-ph] : 1211.3560.
[22] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin, and W. K. Wootters, Phys. Rev. Lett. **76**, 722 (1996); C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K Wootters, Phys. Rev. A **54**, 3824 (1996).
[23] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information (Cambridge Univ. Press, New

York, 2000).

[24] D. D. Dukaric and S. Wolf, arXiv:0808.3317; M. Forster, S. Winkler, and S. Wolf, Phys. Rev. Lett. **102**, 120401 (2009).

[25] See the Appendix.

[26] E. Schrodinger, Proc. Cambridge Philos. Soc. **31**, 553 (1935); **32**, 446 (1936).

[27] H. M. Wiseman, S. J. Jones and A. C. Doherty, Phys. Rev. Lett. **98**, 140402 (2007); S. J. Jones, H. M. Wiseman, and A. C. Doherty, Phys. Rev. A **76**, 052116 (2007).

[28] E. G. Cavalcanti, S. J. Jones, H. M. Wiseman, M. D. Reid, Phys. Rev. A **80**, 032112 (2009).

[29] R. Colbeck, R. Renner, Nature Communications **2**, 411 (2011); R. Colbeck, R. Renner, Phys. Rev. Lett. **108**, 150402 (2012).

[30] S. Kochen and E. P. Specker, J. Math. Mech. **17**, 59 (1967); J. Bub, Interpreting the Quantum World (Cambridge University Press, 1999); A. Peres, J. Phys. A: Math. Gen. **24**, L175 (1991); N.D. Mermin, Phys. Rev. Lett. **65**, 3373 (1990); A. Cabello, Nature (London) **474**, 456 (2011).

[31] J. Bub, arXiv [quant-ph]:1210.6371.